

UR-1528

ER/40685/916

May 1998

Standard-Model-like Chiral Spectra and the Origin of the Families**Otto C. W. Kong**

*Department of Physics and Astronomy,
University of Rochester, Rochester NY 14627-0171.*

Abstract

We outline our approach to understand the family structure through the idea of a SM-like chiral fermion spectrum, one that is derivable from anomaly cancellation conditions in the same way as the one family SM, under an extended symmetry, which after breaking to the SM symmetry yields the three families as the residual chiral content. An example of a relatively simple scalar sector which gives an acceptable symmetry breaking pattern and naturally hierarchical quark mass matrices is also discussed for a successful $SU(4)_A \otimes SU(3)_C \otimes SU(2)_L \otimes U(1)_X$ model. This is a summary of a seminar given at the 37th IUKT winter school, for the proceedings.

Standard-Model-like Chiral Spectra and the Origin of the Families

Otto C. W. Kong

Department of Physics and Astronomy, University of Rochester, Rochester NY 14627.

Each family of the standard model (SM) fermions composes of a chiral spectrum tightly bounded by gauge anomaly cancellation constraints. However, the three family structure and the hierarchy among the masses is a major puzzle.

The fermions in one SM family can be uniquely derived by assuming one multiplet transforming nontrivially under each component gauge group and requiring the minimal chiral spectrum canceling all the anomalies. We seek to understand the family structure through the idea of a SM-like chiral fermion spectrum, one with the same feature as the one family SM under an extended symmetry, which after breaking to the SM symmetry yields naturally the three families as the residual chiral content. The natural choice of the gauge group is $SU(N) \otimes SU(3) \otimes SU(2) \otimes U(1)$. For instance, start with $(\mathbf{4}, \mathbf{3}, \mathbf{2}, \mathbf{1})$ for $N = 4$, the strategy leads to the spectrum

$$(\mathbf{4}, \mathbf{3}, \mathbf{2}, \mathbf{1}), (\bar{\mathbf{4}}, \bar{\mathbf{3}}, \mathbf{1}, \mathbf{x}), (\bar{\mathbf{4}}, \mathbf{1}, \mathbf{2}, \mathbf{y}), (\bar{\mathbf{4}}, \mathbf{1}, \mathbf{1}, \mathbf{z}), \\ (\mathbf{1}, \bar{\mathbf{3}}, \mathbf{2}, \mathbf{a}), (\mathbf{1}, \bar{\mathbf{3}}, \mathbf{1}, \mathbf{b}), (\mathbf{1}, \bar{\mathbf{3}}, \mathbf{1}, \mathbf{c}), (\mathbf{1}, \mathbf{1}, \mathbf{2}, \mathbf{k}), (\mathbf{1}, \mathbf{1}, \mathbf{1}, \mathbf{s}).$$

Solution for the $U(1)$ charges canceling all anomalies exists but fails to give the correct SM embedding. However, analysis of the potentially successful embeddings of the three families suggests that $SU(3)_C \otimes SU(2)_L \otimes U(1)_Y \subset SU(4)_A \otimes SU(3)_C \otimes SU(2)_L \otimes U(1)_X$ works when the above spectrum is augmented with an anomaly-free $SU(4)_A$ multiplet charged under $U(1)_X$. The resulted models have nontrivial $U(1)_Y$ embeddings. Similar constructions with some other N values are also obtained.

These SM-like chiral models have interesting phenomenological predictions. In the case of a specific model with $N = 4$, we also constructed a Higgs sector giving rise to a natural mass hierarchy $m_t, m_b > m_c > m_s > m_d, m_u$. The scalars multiplets are $\phi_0 = (\bar{\mathbf{4}}, \mathbf{1}, \mathbf{1}, \mathbf{9})$ and ϕ_a ($a = 1$ or 2), in $(\bar{\mathbf{4}}, \mathbf{1}, \mathbf{1}, -\mathbf{3})$, together with $SU(2)_L$ doublets $\Phi = (\mathbf{15}, \mathbf{1}, \mathbf{2}, -\mathbf{6})$. A $C_{ab}\phi_{ai}\phi_b^{\dagger j}\Phi_j^k\Phi_k^{\dagger i}$ mass term with natural VEVs for the ϕ_a 's decouples twelve of the fifteen doublets from the EW-scale. There remain two EW Higgs doublets and an extra doublet of singly- and doubly-charged scalars. FCNC constraints can be easily satisfied and the quark mass hierarchy resulted.

References

- Kong, O.C.W. (1996), Mod. Phys. Lett. A 11, 2547.
Kong, O.C.W. (1997), Phys. Rev. D 55, 383.